

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Starane*

EPA Reg. No. 62719-286

EPA 24(c) Special Local Need Registration SLN ID-990017

For Control of Bedstraw in Grasses Grown for Seed
(For Distribution and Use Only in the State of Idaho)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Starane* herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Starane according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use

Starane* herbicide may be applied as a broadcast postemergence treatment using ground equipment or by air. A second application may be made a minimum of 14 days after the first. Starane may be applied in tank mix combination with labeled rates of other herbicides. Read and follow all label directions, including applicable use directions, precautions and limitations on each product label.

Crop/Site/Commodity: Grasses grown for seed, including bluegrass, tall fescue, and perennial ryegrass.

Target Pest/Problem: Bedstraw (*Galium* species).

Key Weeds Controlled, Application Rates and Application Timing to Crop and Weeds:

Crop Stage of Growth	Application Rate (pt/acre)	Application Timing and Specific Use Directions
Established plantings:	2/3	Timing to weeds: Apply when weeds are actively growing and before weeds are 8 inches tall. Only weeds emerged at the time of treatment will be controlled.
Newly seeded plantings:	2/3	Timing to crop: Apply to established grasses in the spring from the tiller stage prior to early boot stage. New plantings may be treated from the 2 true leaf stage of growth prior to early boot stage. Do not apply during boot, flowering, or seed development stage of growth.

Restrictions:

- Do not allow livestock to graze treated areas or harvest treated grasses for forage.
- Do not make more than two applications per growing season.
- **Special Crop Use Restrictions:** Use of Starane according to this labeling is deemed a non-food use, and as such is not for use on fields producing grass for livestock feed. **Do not** feed or graze grass following application of Starane. **Do not** cut treated grass for forage, hay or silage. No portion of the treated field may be used for human or animal feed.

- **Maximum Application Rate:** Do not apply more than 1 1/3 pints per acre of Starane per growing season.
- **Plant-back Restriction:** If replanting is required, only wheat, barley or oats may be planted in treated fields within 120 days following application of Starane.
- **Chemigation:** Do not apply this product through any type of irrigation system.

*Trademark of Dow AgroSciences LLC

Revisions:

D06-075-XXX

Accepted: 06-21-99

Replaces D06-049-012.